

Seeking



in Women's Health

National Women's Annual Clinical
Report 2013



is an art won by training and habituation. We do not act rightly because we have virtue or excellence, but we rather have those because we have acted rightly. We are what we repeatedly do. Excellence, then, is not an act but a habit.

Aristotle (384 BC-322 BC) Greek philosopher.



in Women's Health

Department of Health and Human Services (1996 and 1999).

- comprehensive
- women-friendly
- women-focused
- women-relevant
- integrated
- multidisciplinary

Success dependent upon dual importance of:

- multidisciplinary programs of clinical care
- education of the future providers of women's healthcare

J Womens Health Gend Based Med. 2001 Sep;10(7):627-36.Academic models of clinical care for women: the National Centers of Excellence in Women's. Health.Milliken N, Freund K, Pregler J, Reed S, Carlson K, Derman R, Zerr A, Battistini M, Bowman S, Magnus JH, Sarto GE, Chambers JT, McLaughlin M.

Great



Good



OUR DREAM

Excellence in

- Healthcare delivery
- Clinical and quality outcomes
- Training and research

Achieved by

- Partnership around innovative models of care
- Empowerment of women and their healthcare providers
- Optimal facilities, efficiently utilised and appropriately resourced
- Concentration of special expertise

Resulting in

- an economically sustainable Women's Health Service


Our

Excellence in Women's
Health through

Vision

Empowerment and
Partnership

What should we be held accountable for?



Excellence in Women's Health through Empowerment and Partnership



Safe models of care

Evidence based care
Safe Care
Safe Staffing Levels
Great communication
Respectful relationships
among all staff
Explicit Organizational values

23%

Better Quality Care



Women & family Centered Care

Women and family centered
Women led and empowered
Care responsive to women's
needs
Equity of access for all women

17%

**Safe evidence based
care**



Valued & supported workforce

People valued and nurtured
Teams working together
Training and education of
healthcare professionals
Strong mentoring
Ensuring everyone can
reach their full potential

26%

Engaged Workforce



Optimal resources, efficiently utilized

Efficient use of resources
Working productively
Eliminating system waste
Clinical pathways - KPI'
Living within our means

22%

**Economic
sustainability**



Gold standard Outcomes

Excellent health outcomes
Research supported and valued

11%

**Improved Healthcare
Status**

Integrity

*****

Respect

*****

Innovation

*****

Effectiveness

Safe models of care



A pregnant woman is shown from the side, wearing an orange long-sleeved shirt. She is gently holding her pregnant belly with her right hand. The background is a vibrant landscape featuring a field of yellow wildflowers in the foreground, a clear blue sky with soft white clouds in the middle ground, and a bright sun in the upper left corner creating a lens flare effect.

SUPPORTING WOMEN TO BIRTH WELL

Normal Birth

The baby is born head-first, through the vagina, but may include intervention by a health-care professional to support or help the birth. Interventions included:

- Augmentation
- Intermittent auscultation of FHR
- ARM
- pharmacological pain relief but not epidural
- Active management of 3rd stage

Definitions: Society O&G Canada

Birthing Well

- Empowered woman
- Well informed and prepared
- Realistic expectations
- Able to safely achieve the least interventional birth that is possible given her set of physical and emotional circumstances

Four pathways to birth

- according to ACR

1. Induction of labour
2. Elective CS
3. Emergency CS prior to the onset of labour
4. Spontaneous onset of labour

What is the significance of the order?

Nullipara at term 3422


- 40% BMI>25, 18% > 30, 8%>35
- 5.8% smokers
- Rising maternal age (greatest increase in 35-40 age group)
- High deprivation scores in younger women & Maori & Pacific women

Nullipara at term 3422

- **Ethnicity**
  - 47% European
  - 14% Maori and pacific
  - 35% Asian or Indian
- **Cared for by**
  - 71% under private obstetrician or independent LMC
  - Most complex medically and socially receiving care by NW medical & community teams



3422
Nullipara at
term



3422
Nullipara at
term

What are her chances of a normal
birth at National Women's?

1.50 -60%

2.40-50%

3.30-40%

4.20-30%

5.<20%

3422
Nullipara at term

72 (2.1%)
Emergency pre-
labour Caesarean
section

2987 (87%)
Labour


363 (11%)
Elective
Caesarean section

**Maternal request
91 (20%)**



3422
Nullipara at term

2985(87%)
Labour



1702 (57%)
Spontaneous
labour

1285 (43%)
Induced labour

3422
Nullipara at term


2985 (87%)
Labour



3422 (100%)
Nullipara at term

2985 (87%)
Labour

1702 (57%)
Spontaneous
labour



695 (40.8%)
No Epidural

1008 (59.2%)
Epidural

3422 (100%)
Nullipara at term

2985 (87%)
Labour
of NT

1702 (57%)
Spontaneous
labour



3422
Nullipara at term

2985(87%)
Labour

1702 (50%)
Spontaneous
labour

695 (20.3%)
No epidural

606 (88.6%)
SVD
18% of NT

**36% of
nullipara who
spontaneously
laboured**

66 (9.5%)
Operative vaginal
delivery

**65% nullipara
who started
labour at BC**

13 (1.8%)
Caesarean section

3422
Nullipara at term

2985 (87%)
Labour
of NT

1702 (50%)
Spontaneous
labour

695 (20.3%)
No epidural



3422
Nullipara at term

2985(87%)
Labour

1702 (50%)
Spontaneous
labour

695 (20.3%)
No epidural

616 (18%)
Spontaneous
vaginal delivery

524 (85.1%)
No PPH

64 (10.4%)
PPH>500ml

28 (4.6%)
PPH>1000ml

3422
Nullipara at term

2985 (87%)
Labour

1702 (50%)
Spontaneous
labour

695 (20.3%)
No epidural

616 (18%)
Spontaneous
vaginal delivery



3422
Nullipara at term

2985(87%)
Labour

1702 (50%)
Spontaneous
labour

695 (20.3%)
No epidural

616 (18%)
Spontaneous
vaginal delivery

524 (15.3%)
No PPH

46 (7.3%)
Intact perineum

105 (17.1%)
Episiotomy

4 (0.7%)
3rd or 4th degree
tear

3422
Nullipara at term

2985 (87%)
Labour

1702 (50%)
Spontaneous
labour

695 (20.3%)
No epidural

616 (18%)
Spontaneous
vaginal delivery

524 (15.3%)
No PPH



3422
Nullipara at term

2985(87%)
Labour

1702 (50%)
Spontaneous
labour

695 (20.3%)
No epidural

616 (18%)
Spontaneous
vaginal delivery

524 (15.3%)
No PPH

46 (7.3%)
Intact perineum

Normal birth
18%

Uncomplicated
normal birth
15%

Empowered birth
?

3422
Nullipara at term

2985 (87%)
Labour

1702 (50%)
Spontaneous
labour

695 (20.3%)
No epidural

616 (18%)
Spontaneous
vaginal delivery

524 (15.3%)
No PPH

46 (7.3%)
Intact perineum

A close-up of a person's hand holding a bright yellow, curved object, possibly a piece of fruit or a small container, against a background of a clear blue sky and a green field with yellow flowers. The text is overlaid on the right side of the image.

**Is this what you
expected?**

Could we do better?

**APPROPRIATE AND WELL TIMED
INTERVENTIONS**




INDUCTION OF LABOUR

Induction of labour

- Steady increase in rate over last 20yrs (from 21% to 32.3%)
- Greatest predictor of induced labour
  - Care by high risk medical team (43.2%)
  - Care by diabetes team (70.1%)

IOI Comparison Nationally

Percentage of standard primiparae giving birth



Induction for Prolonged Latent Phase

- Similar rates in Multips (15.2%) & Primips (16.4%)
- Predictor of need for intervention
  - Increased CS rate,
  - epidural rate,
  - operative vaginal delivery rates
- What is going on?
  - Is there an underlying labour abnormality?
  - Or does intervention cause labour abnormalities?

Are we intervening appropriately?



PAIN RELIEF

Analgesia use in labouring women

	Epidural	Pethidine	Water
All	60.3%	8.9%	5.9%
Nullipara Spontaneous Labour	58%	12.2%	14.6%
Nullipara Induced Labour	82.6%	15.4%	4.5%
Multip Spontaneous Labour	27.2%	8.4%	5.2%
Multip Induced Labour	57.9%	7.8%	2.5%



B 6.07 - Rate of epidural incidence in vaginal deliveries

Current Comparison


Asklepios had a rate of 47 %

✓ New data for the Half Year

2012 Jul-Dec

6 months

This KPI was highest at Asklepios at 47 % and lowest at Achilles4 at 11 %



Hospitals are ordered from left to right in order of performance on this KPI.

Associations with use of epidural

Nullipara, sp labour, at term, with SVD

	Epidural (1008)	No Epidural(695)
PPH>500ml	20.6%	10.39%
PPH>1000ml	8.3%	4.6%
3 rd /4 th degree tear	8.8%	0.7%
Intact perineum	4.8%	7.3%


No differences in BMI or median birth weight of babies

To what degree is this related to the epidural?

A collection of surgical instruments, including several pairs of forceps and scissors, are laid out on a blue surgical drape. The instruments are metallic and have various handles and tips. The background is a blurred blue drape.


CAESAREAN SECTION

Overall Caesarean Section rates




Major Contributions to CS rates (56.2%)

Singleton cephalic at term




Previous CS, singleton, cephalic, term



Is this a consequence of care giver preference or are women desirous of CS more likely to seek care from a private obstetrician?


VBAC rate by care giver group




298 had TOL – 78% VBAC (60% 2011)

Is this better selection? Is this the impact of VBAC clinic?

Elective CS after 39 weeks



Elective CS at Term by Gestational Grouping



Safe models of care

Safe models of care

Evidence based care

Safe Care

Safe Staffing Levels

Great communication

Respectful relationships
among all staff

Explicit Organizational values

23%

- Do we increase risk of intervention in low risk women by providing care in a high risk delivery unit?
- Could we do better to prepare women for and support them during birth?
- Could we manage pain in ways other than epidural?
- What is the contribution care giver bias makes to outcome?



SAFE MODELS OF CARE

Clinical Governance Structure




SAFE MODELS OF CARE

Women & Family Centered Care



Women's Voice



- *“I was shocked and shaking, powerless to do anything, feeling helpless, exposed bringing shame.”*
- *“Why didn’t they stop when I asked them to?”*
- *“..and was astounded by the general lack of care and support that I received.”*
- *“I felt unsupported as a mother of a newborn, the communication with regards to the plan of care for me was unclear and conflicting most of the time.....and the acute theatre journey was appalling and humiliating.”*

Women & Family Centered Care

Women & family Centered Care

Women and family centered

Women led and empowered

Care responsive to women's needs

Equity of access for all women
17%

- We need to understand women's needs.
- Explore ways to provide care better linked to community.
- Work on broader aspects of care delivery: empathy, respect, empowerment.



WOMEN & FAMILY CENTERED CARE

The image features five blue, pixelated silhouettes of human figures standing in a horizontal line. The first figure on the left is standing alone. The next three figures are connected to each other and to the first figure by holding hands, forming a continuous chain. The fifth figure on the right is standing alone, not touching the others. The entire scene is set against a plain white background.

VALUED AND SUPPORTED WORKFORCE

What could we do more of?

- 
1. Improved communication
  2. Caring culture
  3. Supporting staff
    - Recruit great quality and committed staff and keep them!
    - More recognition of staff doing great job
    - Continual high demands on staff - work life balance
    - Medical group more involved in normal during training
    - Increased registrar teaching and training

Support for difficult decisions

Audit of Caesarean section at full dilatation

218 women identified

Excluded

- category 1 CS (52)
- Care provided by Private obstetrician (55)
- Not FD or record incomplete (3)

Support for difficult decisions

Analysis of 108 records

Obstetrician was present for 66.6%

Unrelated to

- Time of day
- Day of week
- Category of CS

How did this relate to registrar experience level?

Support for difficult decisions

Double instrumental deliveries (34/5125) 0.66%

- Rate of NICU admission increase from 10.6% to 17.7%
- 3/4th degree tear rate increase from 6.6% to 14.7%
- PPH >1000ml increased from 8.9% to 17.7%

In 2/3 SMO or senior registrar/fellow was listed as the operator

What about the other 1/3?

Support for difficult decisions



Valued and Supported Workforce

Valued & supported workforce

People valued and nurtured
Teams working together
Training and education of
healthcare professionals
Strong mentoring
Ensuring everyone can
reach their full potential

26%

- Senior doctor led care
- Strong support and training for junior doctors
- Team work





**VALUED AND SUPPORTED
WORKFORCE**

A close-up photograph of a stethoscope resting on a US dollar bill. The stethoscope is positioned centrally, with its chest piece and tubing clearly visible. The background is a US dollar bill, showing various details like the number '100', the word 'HUNDRED', and the portrait of Benjamin Franklin. The text 'OPTIMAL RESOURCES EFFICIENTLY UTILIZED' is overlaid in orange, bold, sans-serif font in the lower-left quadrant of the image.

**OPTIMAL RESOURCES EFFICIENTLY
UTILIZED**

OECD Health Data 2010






Improvement Projects

- Eras- Maternity services
- Lactation clinic
- Mapping postnatal flow
- Review of inductions process
- Accelerating release to care 96/98
- Iron infusion
- FBC after caesarian

Regional Collaboration



Optimal Resources Efficiently Utilized

Optimal resources, efficiently utilized

Efficient use of resources

Working productively

Eliminating system waste

Clinical pathways - KPI'

Living within our means

22%

- Lots of work in progress
- Great enthusiasm for improving the way we deliver care



**OPTIMAL RESOURCES EFFICIENTLY
UTILIZED**

A photograph of a vault filled with gold bars. The bars are stacked in rows, creating a strong sense of depth and repetition. The lighting is warm, highlighting the metallic sheen of the gold. The perspective is from a low angle, looking down the length of the stack.

GOLD STANDARD OUTCOMES

NZ Maternity Clinical Indicators

Indicator	All NZ	NW
Standard primip who have a SVD	65.6%	65.1%
Standard primip who has instrumental delivery	16%	15.7%
Standard primip who have a CS	17.9%	19.2%
Standard primip who has an induction	4.8%	4.8%
Standard primip with an intact lower genital tract	27.3%	18.2%
Standard primip with episiotomy no 3/4 th deg tear	22.1%	32.4%
Standard primip with out episiotomy & 3/4 th degree tear	3.2%	2.3%
Standard primip with episiotomy & 3/4 th degree tear	1.3%	1.4%
Women having GA CS	8.3%	6.0%
Women requiring a blood transfusion & CS	3.3%	3.8%
Women requiring a blood transfusion with vaginal birth	1.8%	2.3%
Premature births between 32 & 36 wks	6.7%	7.3%

Maternal Mortality & Morbidity

NZ wide

- 35% of maternal deaths avoidable
- Contributory factors in 55%
  - Organization and management
    - Protocols and guidelines
    - Screening for maternal mental health
  - Personnel or barriers in access to care
- National Women's
  - Early warning score
  - Massive transfusion protocol

AMOSS Notifiable Conditions

Table 76: New Zealand and Australasian rates (per 10,000 maternities) of AMOSS notifiable conditions 2010–2011

	Data collection period	NZ registered births	NZ cases		NZ rate		Australasian rate/ratio ²	95% CI ²
		n	n	NW		NW		
Amniotic fluid embolism	2010–2011	127,728	7	3	0.6	0.4	0.4	0.3–0.6
Antenatal pulmonary embolism	2010–2011	127,728	9	2	0.7	0.27	1.2	1.0–1.4
Eclampsia	2010–2011	127,728	25	2	2	0.27	2.2	1.8–2.6
Influenza with intensive care admission	2010	65,124	8	?	1.2	?	1.9	1.4–2.8
Placenta accreta	2010–2011	127,728	50 ¹	10	3.9	1.33	4.2	3.8–4.8
Peripartum hysterectomy	2010–2011	127,728	56 ¹	12	4.4	1.6	6	5.5–6.5
BMI >50	2010	65,124	297		45.6		26.8	24.9–28.7


1 Twenty-five women had placenta accreta and peripartum hysterectomy.

2 Ellwood D. 2013. Reducing the Impact of PPH. Health Round Table presentation, Sydney.

Perinatal Related Death Rates


District health board (DHB) of residence

Figure 27: Perinatal related death rates (per 1000 births) by DHB of residence (mother) compared to New Zealand perinatal related mortality (with 95% CIs) 2007–2011



Neonatal Encephalopathy Rates

Figure 49: Neonatal encephalopathy rates¹ (per 1000 term births) by DHB of maternal residence (with 95% CIs) compared to New Zealand neonatal encephalopathy rate 2010–2011



¹ No estimates are given for DHBs where number of cases reported was fewer than three.

Pre-term outcomes

- Survival rates from 26 weeks approach those of term babies
- Survival of 24 -27 week babies exceeds that of ANZNN

Gold Standard Outcomes

Gold standard Outcomes

Excellent health outcomes

Research supported
and valued

11%

- Benchmark well
Nationally and
Internationally
- Active research and
audit



**GOLD STANDARD
OUTCOMES**



DEDICATION AND WORK OF THE INTELLIGENCE TEAM

Sheila Stubbs,
Author
Birthing the easy way

“The midwife considers the miracle of childbirth as normal, and leaves it alone unless there's trouble.

The obstetrician normally sees childbirth as trouble; if he leaves it alone, it's a miracle.”



Challenges from Annual clinical Report Presentation 2013

Dr Peter van de Weijer challenged National Women's with regard to the:

- High iatrogenic intervention rates
- High discrepancy in care between care providers
- High iatrogenic complication rates

High Iatrogenic Intervention Rates

	2012		2011	2010
	Mothers	7695	7523	7709
	n	%	%	%
Spontaneous onset of labour	3666	47.6	48.2	52.0
Iatrogenic onset of birth	4029	52.4	51.8	48.0
Elective cesarean section	1278	16.6	15.7	15.9
Emergency cesarean section before labour	268	3.5	3.3	3.5
Induction of labour	2483	32.3	32.7	28.7

Cesarean Section Rate

	2011	2012
Overall C/S rate	32.5%	33.4%
Elective C/S rate	15.7%	16.6%
Emergency C/S rate	16.8%	16.8%
Nullipara elective C/S rate	10.0%	10.8%
Multipara elective C/S rate	20.8%	22.2%
Nullipara emergency C/S rate	24.6%	23.3%
Multipara emergency C/S rate	9.8%	10.5%

The challenges.....

The first challenge is to agree what normal birth is and to collect appropriate data to enable us to report our normal birth rate in the 2013 ACR.

In 2012 **54.2%** women had a **spontaneous vertex delivery** compared to **55.6%** during 2011.

What interventions are included in a normal (or physiological) birth?

- Induction?
- Artificial rupture of membranes in labour?
- Augmentation?
- Episiotomy?
- Post partum haemorrhage?
- Physiological third stage?

The Challenges of Primary Birthing

To reduce iatrogenic interventions it is essential that we increase primary birthing in Auckland - yet the numbers of births are decreasing , **451** women in 2011 reducing to **398** in 2012.

The challenge is to provide real choice to women whilst maintaining safety and confidence in our maternity services

A decorative Maori pattern, likely a haka or koru, is visible on the left side of the slide, rendered in a light blue color.

How many women who live in Auckland
would be able to have a similar birth to this
if they choose to do so?



How many women who live in Auckland
could have an elective C/S if they choose
to do so?

The Challenges of Primary Birthing (2)

- Women's choice
- Respecting and supporting each other
- The media
- Agreeing the 'risk factors' and sticking to the agreement
- The difficulties interpreting the interface between primary and secondary birthing
- Providing normal birth environments in secondary units

Normal Birth Environments St Mary's Birth Centre



Normal Birth Environments






ACR 2013 M O'Brien

Intrapartum



- No of births: 604
- Primiparous: 302 (50%)
- Multiparous: 302 (50%)
- Waterbirths: 188 (31%)
- Women using pool for labour and birth: 302 (50%)



Water Birth




Perineal trauma



- Intact
- 1st degree
- 2nd degree
- 3rd degree
- Episiotomy

Intrapartum transfers: 143 (23.6%)



- prolonged 1st stage
- prolonged 2nd stage
- meconium
- abnormal FHR
- analgesia
- breech
- pv bleeding
- other

The challenges for Secondary Birthing Units

- Providing environments that will enable as 'normal' a birth as possible if that is the woman's choice
- To provide water as an option for pain relief (using telemetry monitoring)
- Not to use interventions unless they are necessary

The Way Forward .....

- Define 'normal' or physiological birth
- Report in 2013 ACR
- Identify normal birth champions
- Change birth environment in DU
- Build on the success of PBAC
- Introduce normal birth pathway
- Identify the barriers to achieving normal birth