

Oral Intake and Gastric Acid Reduction Prophylaxis in Labour

Unique Identifier	NMP200/SSM/093 - v03.00
Document Type	Clinical Guideline
Risk of non-compliance	may result in significant harm to the patient/Te Whatu Ora
Function	Clinical Practice, Patient Care
User Group(s)	Te Toka Tumai Auckland only
• Organisation(s)	Te Whatu Ora Te Toka Tumai Auckland
• Directorate(s)	Women's Health
• Department(s)	Maternity
• Used for which patients?	All maternity women
• Used by which staff?	All clinicians in maternity including access holder lead maternity carers (LMCs)
• Excluded	
Keywords	Antacid prophylaxis, gastric acid prophylaxis, oral intake in labour, fasting guidelines in labouring women, obstetric
Author	Anaesthetist
Authorisation	
• Owner	Service Clinical Director - Secondary Maternity Services
• Delegate / Issuer	Service Clinical Director - Women's Health Anaesthesia
Edited by	Document Control
First issued	18 January 2017
This version issued	19 January 2023 - updated
Review frequency	3 yearly

Contents

1. Purpose of guideline	2
2. Guideline management principles	2
3. Speed of onset of drugs used to reduce gastric acidity	2
4. Indication for acid reduction prophylaxis	2
5. Defining high-risk groups for operative intervention in labour	3
5.1 Maternal risk factors.....	3
5.2 Fetal risk factors	3
6. Treatment for women in labour or for operative intervention	3
6.1 Elective Caesarean sections.....	3
6.2 Lower risk labour	3
6.3 Higher risk labour.....	4
6.4 Decision for operative delivery/trial of forceps/ERPOC/perineal tear repairs/EUA.....	4
6.5 Preparation and administration of intravenous omeprazole injection.....	4
7. Monitoring and audit	4
8. Supporting evidence	5
9. Associated documents.....	5
10. Disclaimer.....	5
11. Corrections and amendments	6

1. Purpose of guideline

Obstetric patients are considered at increased risk of aspiration of gastric contents during general anaesthesia when laryngeal reflexes are reduced. This is associated with significant morbidity and mortality. Obstetric patients are at a higher risk compared to the non-pregnant population because high levels of progesterone cause relaxation of the musculature at the gastro-oesophageal junction and increased gastric residual volume. In addition, there is higher intra-gastric pressure due to the gravid uterus, which causes gastric contents to be forced upwards. Studies have shown that the administration of parenteral opioids such as morphine in labour is associated with delayed gastric emptying.

The risk may be minimised by appropriate acid reduction prophylaxis and restriction of oral intake around the time of delivery for those women who are at a higher risk of operative intervention in the peri-partum period, or those who are undergoing elective Caesarean section.

Acid reduction prophylaxis decreases gastric acid secretion and may decrease gastric volume. It does not affect the pH of the gastric contents that may already be present in the stomach, and therefore must be given some time in advance of general anaesthesia in order to be fully effective. Sodium citrate is a solution that is given orally to neutralise stomach acid just prior to induction of general anaesthesia.

Timely administration of acid reduction prophylaxis and sodium citrate reduces the degree of pulmonary damage should aspiration occur; however, neither will reduce the risk of aspiration or pulmonary damage associated with aspiration of solid food.

2. Guideline management principles

Gastric acid reduction prophylaxis and dietary restriction should be given to those women who are considered higher risk for operative intervention. These women should be identified in labour and offered acid reduction prophylaxis, which should be administered regularly until delivery of the placenta, **after which it should be discontinued.**

These women should only be permitted clear fluids or isotonic sports drinks (e.g. Gatorade or Powerade) and should be discouraged from eating. This is to reduce the risk of aspiration of solid gastric contents if they require operative intervention under general anaesthesia.

3. Speed of onset of drugs used to reduce gastric acidity

Effectiveness of drugs to reduce gastric acidity is dependent on administration timing. Oral omeprazole needs up to three hours to reach maximum effect. Intravenous omeprazole can become effective from 45 to 60 minutes after administration. Sodium citrate is effective immediately, but its duration of action is limited by rate of new gastric acid secretion and speed of gastric emptying.

4. Indication for acid reduction prophylaxis

Omeprazole should be given to those women who are considered higher risk of operative intervention in the peri-partum period. In addition, on making a decision for surgical intervention in labour an appropriate dose and route of administration of intravenous acid reduction therapy should be prescribed by the surgeon if none has been administered during labour.

If a general anaesthetic is considered, sodium citrate should be given orally following discussion with the duty anaesthetist.

Low-risk labouring women do not require acid reduction prophylaxis. However, it should be noted that as labour progresses, and circumstances change, the course of labour may change to a labour with a higher risk of operative intervention.

5. Defining high-risk groups for operative intervention in labour

There is no national or international method for defining women at higher risk of operative intervention in the peri-partum period; however, the following groups of women may be considered at higher risk of operative intervention or general anaesthesia. Use clinical experience and knowledge in identifying women who may require acid reduction prophylaxis as the following list is not exhaustive.

Women with higher risk of operative intervention or general anaesthesia in the peri-partum period:

5.1 Maternal risk factors

- Contra-indication to regional anaesthesia (e.g. coagulation problems, maternal sepsis)
- Previous Caesarean section/vaginal birth after caesarean (VBAC)
- BMI >40 at booking
- Diabetes
- Medical condition of mother, e.g. cardiac disease, respiratory disease
- Previous post-partum haemorrhage >1 L
- Parity ≥5
- Polyhydramnios
- Pre-eclampsia/eclampsia/gestational hypertension
- Slow progress in labour
- Age > 40 years

5.2 Fetal risk factors

- Multiple pregnancy
- Large for gestational age baby (EFW >4.5 kg)
- IUGR
- Pre-term labour
- Signs of fetal distress or meconium

6. Treatment for women in labour or for operative intervention

6.1 Elective Caesarean sections

- Nil by mouth: six hours for food and two hours for clear fluids, including the pre-operative Nutricia carbohydrate drink prior to their surgery
- Omeprazole 20 mg capsule on arrival in ORDA
- Those women who are having their surgery under general anaesthetic should also receive oral sodium citrate (30 mL of 300 mmol/L). This should be given within 15 minutes of induction of general anaesthesia

6.2 Lower risk labour

- No gastric acid reduction prophylaxis required

6.3 Higher risk labour

At start of labour, or if a previously lower risk labour develops risk factors for operative intervention:

- Woman should be advised to have clear fluids orally only (no food)
- Gastric acid reduction prophylaxis should be prescribed on the ONCE ONLY section of the National Medical Chart and given as:
 - Omeprazole 20 mg orally (one dose lasts up to 24 hours)

6.4 Decision for operative delivery/trial of forceps/ERPOC/perineal tear repairs/EUA

- A single dose of intravenous omeprazole (40 mg) injection should be given once the decision has been made for surgical intervention if omeprazole has not already been given in the previous 24 hours. Prescribe this on the ONCE ONLY section of the National Medical Chart.
- Oral omeprazole takes at least two hours to be effective, and the intravenous injection preparation takes around 60 minutes to have clinical effect. Therefore, these should be administered as soon as possible after decision for theatre has been made, unless clinical urgency dictates otherwise.
- If a general anaesthetic is considered, then sodium citrate (30 mL of 300 mmol/L) should be given orally. This is normally administered by the duty anaesthetist in the hand-over or operating room.

The administration of omeprazole or sodium citrate **SHOULD NOT** delay transfer to theatre.

6.5 Preparation and administration of intravenous omeprazole injection

Follow these steps.

Step	Action
1.	Use omeprazole injection with diluent provided. DO NOT use sodium chloride 0.9%
2.	Reconstitute 40 mg vial with the 10 mL diluent provided (final concentration 4 mg/mL)
3.	Draw up 10 mL of diluent provided, no other solution must be added or used
4.	Slowly add approximately 5 mL of the diluent to the vial of powder
5.	Withdraw as much air as possible from the vial to reduce the positive pressure
6.	Transfer the remaining diluent into the vial
7.	Rotate and shake the vial to dissolve the powder
8.	Prepare immediately before use (or discard)
9.	Discolouration may occur if incorrect reconstitution technique is used – do not use if solution is discoloured or if any precipitate is present
10.	Administration: <ul style="list-style-type: none">• Flush with sodium chloride 0.9% before and after administration• Inject slowly over at least 2.5 to 5 minutes, at a rate not exceeding 4 mL/minute

7. Monitoring and audit

Clinical audit should be undertaken to review guideline compliance; 100% compliance (excluding exempted women) is expected for both dietary restriction and antacid prophylaxis in all women presenting with a higher risk of operative delivery or anaesthetic intervention in the peri-partum period. Gastric acid reduction prophylaxis should be prescribed according to women's individual needs. Exceptions to this include women who decline pharmacological gastric acid reduction prophylaxis, allergy to the drugs or other contraindication.

8. Supporting evidence

- Mendelson, C. L. (1946). The aspiration of stomach contents into the lungs during obstetric anesthesia. *Obstetrical & Gynecological Survey*, 1(6), 837-839.
- Clark, K., Lam, L. T., Gibson, S., & Currow, D. (2009). The effect of ranitidine versus proton pump inhibitors on gastric secretions: a meta-analysis of randomised control trials. *Anaesthesia*, 64(6), 652-657.
- Stuart, J. C., Kan, A. F., Rowbottom, S. J., Yau, G., & Gin, T. (1996). Acid aspiration prophylaxis for emergency Caesarean section. *Anaesthesia*, 51(5), 415-421.
- Lewis, G. (2004). *Confidential Enquiry into Maternal and Child Health: Why mothers die 2000-2002*. London: RCOG Press, 122-134.
- Gyte, G. M., & Richens, Y. (2010). Routine prophylactic drugs in normal labour for reducing gastric aspiration and its effects. *Cochrane Database of Systematic Reviews*, (3).
- National Institute for Health and Care Excellence. (2014). Intrapartum for healthy women and babies. Retrieved from: <https://www.nice.org.uk/guidance/cg190>.
- Levy, D. M. (2006). Pre-operative fasting—60 years on from Mendelson. Continuing Education in Anaesthesia, *Critical Care & Pain*, 6(6), 215-218.
- Medsafe. (2019). *New Zealand Data Sheet for Omeprazole tablets (Losec)*. Retrieved from: <https://www.medsafe.govt.nz/profs/datasheet/l/Loseccap.pdf>.
- Medsafe. (2019). *New Zealand Data Sheet for Intravenous Omeprazole (Dr. Reddy's)*. Retrieved from: <https://www.medsafe.govt.nz/profs/Datasheet/o/Omeprazoleinj.pdf>
- Paranjothy, S., Griffiths, J. D., Broughton, H. K., Gyte, G. M., Brown, H. C., & Thomas, J. (2014). Interventions at caesarean section for reducing the risk of aspiration pneumonitis. *The Cochrane Library*.
- American Society of Anesthesiologists Task Force on Obstetric Anesthesia. (2007). Practice guidelines for obstetric anesthesia: an updated report by the American Society of Anesthesiologists Task Force on Obstetric Anesthesia. *Anesthesiology*, 106(4), 843.
- Singata, M., Tranmer, J., & Gyte, G. M. (2010). Restricting oral fluids and food intake during labour *Cochrane Database of Systematic Reviews*.
- Whittington, R. M., Robinson, J., & Thompson, J. (1979). Fatal aspiration (Mendelson's) syndrome despite antacids and cricoid pressure. *The Lancet*, 314(8136), 228-230.

9. Associated documents

- Caesarean Section (CS) - Pre, Peri & Post-Op Care
- Caesarean Section - Acute - Level 8 Operating Room
- Caesarean Section (CS) - Post Anaesthesia Care Unit (PACU)
- Perineal Tears – Third and Fourth Degree
- Intrapartum Care – Physiological Labour and Birth
- Surgical Safety Checklist
- Medications - Prescribing

Clinical forms

- Preoperative Patient Assessment and Check in CR4135

10. Disclaimer

No guideline can cover all variations required for specific circumstances. It is the responsibility of the health care practitioners using this Te Toka Tumai Auckland guideline to adapt it for safe use

If printed, this document is only valid for the day of printing.

within their own institution, recognise the need for specialist help, and call for it without delay, when an individual patient falls outside of the boundaries of this guideline.

11. Corrections and amendments

The next scheduled review of this document is as per the document classification table (page 1). However, if the reader notices any errors or believes that the document should be reviewed **before** the scheduled date, they should contact the owner or [Document Control](#) without delay.