

Bladder Care Postpartum and Management of Urinary Retention

| | |
|----------------------------|--|
| Unique Identifier | NMP200/SSM/073 – v05.00 |
| Document Type | Clinical Guideline |
| Risk of non-compliance | very unlikely to result in harm to the patient/DHB |
| Function | Clinical Practice, Patient Care |
| User Group(s) | ADHB only |
| • Organisation(s) | Auckland District Health Board (ADHB) |
| • Directorate(s) | National Women’s Health |
| • Department(s) | Maternity |
| • Used for which patients? | Postpartum maternity women |
| • Used by which staff? | All clinicians in maternity including access holder lead maternity carers (LMCs) |
| • Excluded | |
| Keywords | n/a |
| Author | Midwifery Educator - Physiotherapist and Urogynaecologist |
| Authorisation | |
| • Owner | Service Clinical Director - Secondary Maternity Services |
| • Delegate / Issuer | |
| Edited by | Clinical Policy Advisor |
| First issued | April 2008 |
| This version issued | 20 November 2017 - reviewed |
| Review frequency | 3 yearly |

Contents

| | |
|---|---|
| 1. Purpose of guideline | 2 |
| 2. Guideline management principles and goals | 2 |
| 3. Management | 2 |
| 3.1 Overt retention..... | 4 |
| 3.2 Covert retention | 5 |
| 4. Admission assessment..... | 6 |
| 5. First 6 hours post-delivery or removal of IDC assessment | 6 |
| 6. Urinary retention | 7 |
| 7. Removal of catheter (trial of void) management | 8 |
| 8. Supporting evidence | 8 |
| 9. Associated Auckland DHB documents..... | 9 |
| 10. Disclaimer | 9 |
| 11. Corrections and amendments | 9 |

1. Purpose of guideline

The purpose of this guideline is to assist health professionals in bladder care during the postpartum period, with the aim of preventing urinary retention and its long-term consequences within Auckland District Health Board (Auckland DHB).

2. Guideline management principles and goals

Hormone induced reduction in smooth muscle tone decreases bladder tone (hypotonia) during pregnancy and for a period following birth. These changes may persist for days or longer in some women with the risk of over distension of the postpartum bladder (Saultz, 1991, see supporting evidence section).

Vigilant surveillance of bladder function and early intervention where problems exist should prevent permanent bladder damage and long-term voiding problems (Rizvi, 2005, see supporting evidence section).

While all women in the immediate postpartum period have the potential to experience urinary problems, several factors increase the risk:

- Prolonged/difficult labour
- Delay in the second stage
- Assisted birth
- Caesarian birth
- Epidural analgesia, particularly with local anesthetic
- Perineal/vulval trauma
- Over distension of the bladder during/immediately following birth
- Large infant > 4 kg
- English as a second language
- Pain
- Constipation

Aims of care

- To assess bladder function
- To detect any deviation/s from normal
- To carry out timely preventative measures to avoid complications of urinary dysfunction following birth


3. Management

There are two types of urinary retention that can affect a woman in the postpartum period:


- Overt retention: Symptomatic inability to void spontaneously within six hours of birth or removal of IDC

- Covert retention: Non symptomatic increased post void residual volumes after birth or removal of IDC

3.1 Overt retention



3.2 Covert retention



4. Admission assessment

The initial bladder assessment should include:

- A review of the labour and birth history to detect any risk factors
- History of urological problems
- Bladder palpation
- Check to see if the woman has voided after vaginal birth
- IDC in situ –check that it is draining

An initial assessment should provide information on:

- The presence of any urinary problems
- Risk factors that may contribute to urinary problems

5. First 6 hours post-delivery or removal of IDC assessment

Continue to assess bladder functioning 2 hourly. If unable to void or quantity or flow is abnormal at 4 hours, refer to the postpartum overt urinary retention flowchart (adapted from the WHA Guidelines, 2009).

Notes:

- During the night: If there is no history of urological problems, use opportunities when the woman is awake to check bladder. A woman with a history may require 2 hourly checking
- Onset and progression of urinary retention may be gradual and asymptomatic
- It can take 8 hours for the bladder to regain sensation following epidural analgesia

Assessment:

- Establish by questioning void or no void
- If yes to void, ask the woman if she is experiencing any discomfort or difficulty when voiding
- Check the frequency with which urine is passed
- Ask volume and quality of flow with each void
- Examine the woman's abdomen for displacement of the uterus and swelling of the lower abdomen
- Palpate the woman's bladder
- Establish by questioning void or no void

The woman may complain of overt symptoms (symptomatic inability to void spontaneously within six hours of birth or removal of IDC):

- An inability to void
- Increasing lower abdominal pain
- Urgency

- Straining to void
- Involuntary loss of urine
- Voiding frequent small amounts (retention with overflow)

Note: A distended bladder displaces the uterus upward and to the right side. There may also be a painful cystic swelling palpable in the suprapubic region.

If no void at 4 hours either post birth or removal of IDC use supportive measures, such as ambulation, privacy, shower, hands under cold running water, warm flannel over bladder or if necessary appropriate analgesia for pain relief to enhance the likelihood of micturition. Ensure adequate fluid intake and commence Fluid Balance Chart.

Monitor a further 2 hours: (i.e. until 6 hours post delivery or sooner if discomfort)

- If void and volume > 200 mL continue with supportive measures and encourage 2 - 3 hourly voiding
- If no void or volume < 200 mL drain bladder with IDC

When inserting a catheter:

- Use a Foley's catheter
- Use a strict aseptic technique
- Send CSU to lab
- Document on the Fluid Balance Chart

Note: Using a Foley catheter, instead of an in-out catheter prevents the risk of introducing bacteria into the urinary tract from a second catheterization should an indwelling catheter be required.

A woman may have covert urinary retention (non-symptomatic increased post void residual volumes after birth or removal of IDC):

- Ability to void
- But no urge to void
- No obvious symptoms of retention

Refer to postpartum covert urinary retention flowchart (adapted from the WHA Guidelines, 2009).

6. Urinary retention

Alert obstetric team.

Diagnosed by symptoms and volume drained following insertion of IDC.

- Residual urinary volume of 150 - 700 mL will require IDC for 24 hours
- Residual urinary volume > 700 mL will require IDC 48 hours

- Residual volumes of > 1500mls require discussion with the Urogynaecology team

A woman who has a residual volume of more than 700 mL is more likely to require repeat catheterization (Ching-Chung, 2002, see supporting evidence section). After a failed trial of removal of catheter discussion with urogynae team is required for further management and refer to the ward physiotherapist.

Catheterization rests the over distended bladder allowing it to gain its elastic recoil.

It is advisable to remove urinary catheters early in the day to allow time for careful and regular post catheterization bladder assessment.

7. Removal of catheter (trial of void) management

Encourage 2 - 3 hourly voiding and document voids until normal voiding patterns are established and two measured voids of 200 mL or greater are obtained.

Reassess the bladder as documented and follow the appropriate flowchart for postpartum overt urinary retention or postpartum covert urinary retention (adapted from the WHA Guidelines, 2009).

Document all findings on the Fluid Balance Chart and in the clinical record.

Persistent urinary retention and large urinary residuals will require long term resting of the bladder and management by the obstetric team in conjunction with Urogynaecology.

Note: Bladder scanners are not a reliable measurement of residual volumes in the postpartum woman and are not recommended for use. The automatic calculation is rendered inaccurate because of the volume of the involuting uterus and its tendency to distort the bladder outline (Pallis & Wilson 2003, see supporting evidence section).

8. Supporting evidence

- Carley M.E et al., 2002: [Factors that are associated with clinically overt postpartum urinary retention after vaginal delivery](#). American journal of Obstetrics and Gynaecology
- Ching-Chung, L Shuenn-Dhy, C, Ling-Hong, T. 2002 [Postpartum urinary retention: assessment of contributing factors and long term clinical impact](#). Australia and New Zealand Journal and Obstetrics and Gynaecology, 42 (4) 367-370
- R.M. Rizvi, Z.S. Khan and Z. Khan, [Diagnosis and management of postpartum urinary retention](#), Int J Obstet Gynecol 91 (2005) (1), pp. 71–72
- Rogers, R G & leeman, L.I.L. 2007 [Postpartum genitourinary changes](#). Urological Clinics of North America, 34 13-21
- Saultz J.W., William L., Toffler W.L. & Shackles J. Y. (1991) [Post partum urinary retention](#). Journal of the American Board of Family Practitioners 4, 341-344

- Pallis L. M. & Wilson M. (2003) [Ultrasound assessment of bladder volume: Is it valid after delivery?](#) Australian and New Zealand Journal of Obstetrics and Gynaecology 43, 453-456
- Van Os, A.F.M., & Van der Linden, P.J.Q. (2006). [Reliability of an automatic ultrasound system in the postpartum period in measuring urinary retention.](#) Acta Obstetrica et Gynecologica Scandinavica, 85, 604-607
- WHA Clinical Practice Guideline March 2009
- World Health Organisation. [Sexual and reproductive health.](#) Retrieved 30 April 2007
- World Health Organization. [Postpartum care of the mother and newborn: a practical guide. Maternal and newborn health](#)
- Yip, S., Sahota, D., Pang, M., & Chang, A. (2004). [Postpartum urinary retention.](#) Acta Obstetrica et Gynecologica Scandinavica, 83, 887-891
- Yip, S., Sahota, D., Pang, M., & Chang, A. (2005). [Screening test model using duration of labour for the detection of postpartum urinary retention.](#) Neurology and Urodynamics, 24, 248-253
- Zaki, M.M., Panid, M. & Jackson, S. (2004). [National survey for intrapartum bladder care: assessing the need for guidelines.](#) BJOG: an International Journal of Obstetrics and Gynaecology, 111, 874-876

9. Associated Auckland DHB documents

- [Bladder Care Post Gynaecology & Urogynaecology Surgery](#)

10. Disclaimer

No guideline can cover all variations required for specific circumstances. It is the responsibility of the health care practitioners using this Auckland DHB guideline to adapt it for safe use within their own institution, recognise the need for specialist help, and call for it without delay, when an individual patient falls outside of the boundaries of this guideline.

11. Corrections and amendments

The next scheduled review of this document is as per the document classification table (page 1). However, if the reader notices any errors or believes that the document should be reviewed **before** the scheduled date, they should contact the owner or [Clinical Policy Advisor](#) without delay.